

# Virginia Tech Regional Math Contest 1980<sup>1</sup>

## Problem 1

Let  $*$  denote a binary operation on a set  $S$  with the property that

$$(w * x) * (y * z) = w * z \text{ for all } w, x, y, z \in S$$

Show:

- (a) If  $a * b = c$ , then  $c * c = c$ .
- (b) If  $a * b = c$ , then  $a * x = c * x$  for all  $x \in S$ .

## Solution 1

### Part (a)

Use the given equation, substitute, and simplify:

$$\begin{aligned} (w * x) * (y * z) &= w * z && \text{given} \\ (a * b) * (a * b) &= a * b && \text{substitute } w = a, x = b, y = a, z = b \\ c * c &= c && \text{simplify using } a * b = c \end{aligned}$$

This proves that if  $a * b = c$ , then  $c * c = c$ . ■

### Part (b)

Use the given equation with some substitutions and utilize the result from part 1:

$$\begin{aligned} (w * x) * (y * z) &= w * z && \text{given} \\ (c * c) * (x * x) &= c * x && \text{substitute } w = c, x = c, y = x, z = x \\ c * (x * x) &= c * x && \text{simplify using } c * c = c \text{ from part (a)} \\ (a * b) * (x * x) &= c * x && \text{expand using } a * b = c \\ a * x &= c * x && \text{simplify using given equation} \end{aligned}$$

This proves that if  $a * b = c$ , then  $a * x = c * x$  for all  $x \in S$ . ■

## Problem 2

The sum of the first  $n$  terms of the sequence

$$1, (1 + 2), (1 + 2 + 2^2), \dots, (1 + 2 + \dots + 2^{k-1}), \dots$$

is of the form  $2^{n+R} + Sn^2 + Tn + U$  for all  $n > 0$ . Find  $R, S, T$  and  $U$ .

## Solution 2

The given sequence is already defined for us in the general case in the last element:

$$1, (1 + 2), (1 + 2 + 2^2), \dots, (1 + 2 + \dots + 2^{k-1}), \dots$$

The  $i$ th term of this sequence can thus be written as follows:

$$a_i = \sum_{i=1}^n 2^{i-1}$$

Then, the sum of the first  $n$  terms in the sequence is:

---

<sup>1</sup>VTRMC problems source: <https://personal.math.vt.edu/plinnell/Vtregional/>

$$\sum_{i=1}^n a_i = \sum_{i=1}^n \sum_{k=1}^i 2^{k-1}$$

However, we don't actually have to expand or simplify the double-summation. Since we have 4 unknowns, we can just compute the first 4 values of  $A_n$ , set them equal, and solve a system of 4 equations with 4 unknowns.

Let  $A_n = \sum_{i=1}^n a_i$ , and let's compute several of the initial values:

$$\begin{aligned} A_1 &= a_1 & &= 1 & & & &= 1 \\ A_2 &= a_1 + a_2 & &= 1 + (1 + 2) & & & &= 4 \\ A_3 &= a_1 + a_2 + a_3 & &= 1 + (1 + 2) + (1 + 2 + 4) & & & &= 11 \\ A_4 &= a_1 + a_2 + a_3 + a_4 & &= 1 + (1 + 2) + (1 + 2 + 4) + (1 + 2 + 4 + 8) & & & &= 26 \end{aligned}$$

Additionally, per the given problem statement, we also know that

$$A_n = 2^{n+R} + Sn^2 + Tn + U$$

so let's set them equal for  $n = 1, 2, 3, 4$ :

$$A_1 = 1 = 2^{1+R} + S + T + U \quad (n = 1) \quad (1)$$

$$A_2 = 4 = 2^{2+R} + 4S + 2T + U \quad (n = 2) \quad (2)$$

$$A_3 = 11 = 2^{3+R} + 9S + 3T + U \quad (n = 3) \quad (3)$$

$$A_4 = 26 = 2^{4+R} + 16S + 4T + U \quad (n = 4) \quad (4)$$

We now have 4 equations and 4 unknowns.

Reorganizing equation (1) to put  $R$  on one side, and all the other variables on the other side, we get:

$$2^{1+R} + S + T + U = 1 \quad \text{from eq. 1}$$

$$2^{1+R} = 1 - S - T - U \quad \text{solve for } 2^{1+R} \quad (5)$$

since  $2^{1+R}$  appears in all other equations and  $R$  does not appear as a separate variable, we don't have to solve for  $R$ , and can substitute the result in (5) into the remaining equations (2), (3), and (4) to remove the use of variable  $R$ , leaving us with 3 equations and 3 unknowns ( $S, T, U$ ).

Let's now take equation (2), separate out  $2^{1+R}$  and substitute the result from (5):

$$2 \cdot 2^{1+R} + 4S + 2T + U = 4 \quad \text{from eq. 2}$$

$$2 \cdot (1 - S - T - U) + 4S + 2T + U = 4 \quad \text{substitute from eq. 5}$$

$$2 - 2S - 2T - 2U + 4S + 2T + U = 4 \quad \text{expand}$$

$$2 + 2S - U = 4 \quad \text{simplify}$$

$$2S - 2 = U \quad \text{solve for } U \quad (6)$$

Now let's take equation (3) and similarly substitute from both (5) and (6):

$$\begin{aligned}
2^2 \cdot 2^{1+R} + 9S + 3T + U &= 11 && \text{from eq. 3} \\
2^2 \cdot (1 - S - T - U) + 9S + 3T + U &= 11 && \text{substitute } 2^{1+R} \text{ from eq. 5} \\
4 - 4S - 4T - 4U + 9S + 3T + U &= 11 && \text{expand} \\
5S - T - 3U &= 7 && \text{simplify} \\
5S - T - 3(2S - 2) &= 7 && \text{substitute } U \text{ from eq. 6} \\
5S - T - 6S + 6 &= 7 && \text{expand} \\
-S - T &= 1 && \text{simplify} \\
-S - 1 &= T && \text{solve for } T \text{ (7)}
\end{aligned}$$

Finally, let's take equation (4) and substitute the results from (5), (6), and (7) to eliminate all other variables and solve for  $S$ :

$$\begin{aligned}
2^3 \cdot 2^{1+R} + 16S + 4T + U &= 26 && \text{from eq. 4} \\
2^3 \cdot (1 - S - T - U) + 16S + 4T + U &= 26 && \text{substitute } 2^{1+R} \text{ from eq. 5} \\
8 - 8S - 8T - 8U + 16S + 4T + U &= 26 && \text{expand} \\
8 + 8S - 4T - 7U &= 26 && \text{simplify} \\
8 + 8S - 4(-S - 1) - 7U &= 26 && \text{substitute } T \text{ from eq. 7} \\
8 + 8S - 4(-S - 1) - 7(2S - 2) &= 26 && \text{substitute } U \text{ from eq. 6} \\
8 + 8S + 4S + 4 - 14S + 14 &= 26 && \text{expand} \\
(8S + 4S - 14S) + (8 + 4 + 14) &= 26 && \text{combine like terms} \\
-2S + 26 &= 26 && \text{simplify} \\
-2S &= 0 && \text{simplify} \\
S &= 0 && \text{solve for } S \text{ (8)}
\end{aligned}$$

Now that we have  $S = 0$ , we can go back and substitute this into earlier equations to find the other unknowns. Let's start with eq. (7) and compute  $T$ :

$$T = -S - 1 = 0 - 1 = -1$$

Similarly, using eq. (6), we can compute  $U$ :

$$U = 2S - 2 = 0 - 2 = -2$$

and finally, to compute  $R$ , we can use eq. (5):

$$2^{1+R} = 1 - S - T - U = 1 - 0 - (-1) - (-2) = 4 = 2^2$$

Since  $2^{1+R} = 2^2$ , we know that  $R = 1$ . Hence, here are all the unknown values:

$$\boxed{R = 1; S = 0; T = -1; U = -2} \quad \blacksquare$$

### Verification

Now that we've solved for  $R$ ,  $S$ ,  $T$ , and  $U$ , let's verify some of the  $A_n$  values to make sure it works out:

$$\begin{aligned}
A_1 &= \boxed{1} = 2^{1+R} + S + T + U = 2^{1+1} + 0 - 1 - 2 = 4 - 3 = \boxed{1} \\
A_2 &= \boxed{4} = 2^{2+R} + 4S + 2T + U = 2^{2+1} + 0 - 2 - 2 = 8 - 4 = \boxed{4} \\
A_3 &= \boxed{11} = 2^{3+R} + 9S + 3T + U = 2^{3+1} + 0 - 3 - 2 = 16 - 5 = \boxed{11} \\
A_4 &= \boxed{26} = 2^{4+R} + 16S + 4T + U = 2^{4+1} + 0 - 4 - 2 = 32 - 6 = \boxed{26}
\end{aligned}$$

Checks out!